



## PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

### **EROMYCIN 250 TABLET** **EROMYCIN 500 TABLET** Erythromycin 250mg Tablet Erythromycin 500mg Tablet

#### **Read this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again before, during or after use of this medicine.
- If you have any further questions, ask your health care provider.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects become serious, or if you notice any side effects not listed in this leaflet, please inform your health care provider.

In this leaflet:

1. What Eromycin Tablet is and what it is used for
2. What you need to know before you take Eromycin Tablet
3. How to use Eromycin Tablet
4. Possible side effects
5. How to store Eromycin Tablet
6. Contents of the pack and other information

#### **1. WHAT EROMYCIN TABLET IS AND WHAT IT IS USED FOR**

The name of your medicine is Eromycin Tablets.

Eromycin contains the active ingredient erythromycin which belongs to a group of medicines called macrolide antibiotics.

Eromycin Tablets are used to prevent and treat infections such as:

- Throat and sinus infections
- Chest infections, such as bronchitis and pneumonia
- Ear infections
- Mouth and dental infections
- Eye infections
- Skin and tissue infections, such as acne
- Stomach and intestinal infections
- Prevention of infection following burns, operations or dental procedures
- Other infections, such as sexually transmitted diseases, bone infections or scarlet fever

#### **2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE EROMYCIN TABLET**

**You should not receive Eromycin Tablets if you:**

- Have been told that you are allergic to erythromycin or other macrolide antibiotics such as clarithromycin or azithromycin or any of the other ingredients in these tablets.
- Are currently taking a medicine called
  - Ergotamine or dihydroergotamine (used to treat migraines) while taking Eromycin as this may cause serious side effects.
  - Terfenadine or astemizole or mizolastine (widely taken for hayfever and



## PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

### **EROMYCIN 250 TABLET**

### **EROMYCIN 500 TABLET**

Erythromycin 250mg Tablet

Erythromycin 500mg Tablet

allergies), domperidone (for nausea (feeling sick) and vomiting (being sick)), cisapride (for stomach disorders) or pimozide or amisulpride (for psychiatric conditions) while receiving erythromycin, as combining these drugs can sometimes cause serious disturbances in heart rhythm. Consult your doctor for advice on alternative medicines you can take instead.

- Simvastatin (used to lower cholesterol and triglycerides (types of fat) in the blood).
- Tolterodine (used for treating overactive bladder with symptoms of urinary frequency, urgency, and leakage)
- Have abnormally low levels of potassium or magnesium in your blood (hypomagnesaemia or hypokalaemia).
- or someone in your family has a history of heart rhythm disorders (ventricular cardiac arrhythmia or torsades de pointes) or an abnormality of the electrocardiogram (electrical recording of the heart) called “long qt syndrome”.
- Are taking colchicine (used for treatment of gout and arthritis) whilst taking erythromycin as this may cause serious side effects.
- Have any liver problems or have been told that any drugs you are taking can cause liver problems
- Have previously experienced diarrhoea following the use of antibiotics;
- Are pregnant and have been told that you have a sexually transmitted disease called syphilis. In this case erythromycin may not be effective for preventing the transfer of this infection to your baby. Consult your doctor before receiving erythromycin. Alternatively, if you were treated for early stages of syphilis during your pregnancy, and your child is under 1 year and is prescribed erythromycin, consult your doctor before giving erythromycin to your child.
- Are treating a young child with antibiotics and they are irritable or vomit when fed, you should contact your physician immediately.
- Suffer from a condition called myasthenia gravis, which causes muscle weakness, consult your doctor before receiving erythromycin.
- Are taking erythromycin with ‘statins’ such as simvastatin or lovastatin (used to lower cholesterol) as serious side effects can occur.
- You are taking other medicines which are known to cause serious disturbances in heart rhythm.
- If you have heart problems.

If any of these apply to you, or if you are not sure, tell your doctor.

#### **Taking other medicines**

Please tell your doctor if you are taking, or have recently taken any other medicines,



## PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

### **EROMYCIN 250 TABLET**

### **EROMYCIN 500 TABLET**

Erythromycin 250mg Tablet

Erythromycin 500mg Tablet

including any medicines obtained without a prescription.

This is especially important if you are taking medicines from the following families:

- Astemizole, terfenadine or mizolastine (used to treat allergies such as hayfever);
- Domperidone (used to treat nausea (feeling sick) and vomiting (being sick));
- Pimozide (used to treat mental problems);
- Ergotamine or dihydroergotamine (used to relieve migraine);
- Cisapride (used to treat stomach disorders);
- Statins (used to help lower cholesterol levels e.g. Lovastatin and simvastatin);
- Protease inhibitors (used to treat viral infections e.g. Saquinavir);
- Oral contraceptives.

#### **This is also important if you are taking medicines called:**

- Colchicine (used to treat gout and arthritis)
- Cimetidine and omeprazole (used to treat acid reflux and other related conditions)
- Clarithromycin, rifabutin, or rifampicin (medicines used to treat different types of bacterial infection)
- Fluconazole, ketoconazole and itraconazole (medicines used to treat fungal infections)
- Digoxin, quinidine or disopyramide (used to treat heart problems)
- Cilostazol (a medicine used to treat peripheral circulation problems)
- Hexobarbitone, phenobarbital or midazolam (used as sedatives)
- Anticoagulants e.g. Warfarin, acenocoumarol and rivaroxaban (used to thin the blood);
- Valproate, carbamazepine or phenytoin (used to control epilepsy)
- Theophylline (used to treat asthma and other breathing problems)
- Ciclosporin or tacrolimus (used following organ transplants)
- Bromocriptine (used to treat parkinson's disease)
- Zopiclone or triazolam/alprazolam (used to help you sleep or relieve states of anxiety)
- Alfentanil (a medicine used to provide pain relief)
- Methylprednisolone (used to help suppress the body's immune system – this is useful in treating a wide range of conditions)
- St john's wort (a herbal medicine used to treat depression)
- Verapamil (used to treat high blood pressure and chest pain)
- Vinblastine (used to treat certain types of cancer)
- Sildenafil (used to treat erectile dysfunction).

If you or your child goes for any medical tests, tell your doctor that you are taking Erythromycin, as this may interfere with some test results.



## PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

### **EROMYCIN 250 TABLET**

### **EROMYCIN 500 TABLET**

Erythromycin 250mg Tablet

Erythromycin 500mg Tablet

#### **Pregnancy and breast-feeding**

Eromycin should be used by women during pregnancy only if clearly needed.

If you are pregnant or think you may be pregnant, or if you are breast-feeding, please consult your doctor before taking this medicine.

### **3. HOW TO TAKE EROMYCIN TABLETS**

Always take your medicine exactly as instructed. Ask your doctor or pharmacist if you are not sure.

Always take Eromycin Tablets exactly as your doctor has told you. You should check with your doctor if you are not sure.

The usual dose of Eromycin Tablets for adults and children over 8 years is:

1-2 g daily in divided doses i.e., four to eight 250 mg tablets daily or two to four 500 mg tablets daily, taken just before or with meals or food. If you have a bad infection, you may be told to take up to 4 g daily.

Continue to take this medicine until the course is completed or until your doctor tells you to stop; do not stop taking your medicine, even if you feel better. If you stop the treatment too early your problem could come back.

#### **If you take more Eromycin Tablet than you should**

If you accidentally take more medicine in one day than your doctor has told you to, or if a child has taken some of the medicine by mistake, contact your doctor or go to your nearest hospital emergency department immediately.

An overdose of Eromycin could cause temporary hearing loss, nausea, vomiting and diarrhoea.

#### **If you forget to take Eromycin Tablet**

If you forget to take a dose of your medicine, take it as soon as you remember. Do not take more medicine in one day than your doctor tells you to.

If you have any further questions on the use of this product ask your doctor or pharmacist

### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Eromycin Tablets can cause side effects, although not everybody gets them.



## PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

### **EROMYCIN 250 TABLET**

### **EROMYCIN 500 TABLET**

Erythromycin 250mg Tablet

Erythromycin 500mg Tablet

If you notice any of the following, contact your doctor immediately:

- Difficulty breathing
- Fainting
- Swelling of the face, lips or throat
- Skin rashes
- Severe skin reactions including large fluid-filled blisters, sores and ulcers
- Ulcers in the mouth and throat, as these may be signs of an allergic reaction.

Contact a doctor immediately if you experience a serious skin reaction: a red, scaly rash with bumps under the skin and blisters (exanthematous pustulosis). The frequency of this side effect is not known (cannot be estimated from the available data).

#### **Other side effects of Eromycin include:**

- Diarrhoea which may be severe or prolonged and may contain blood or mucus
- Feeling sick or being sick
- Increase in a particular type of white blood cells (eosinophilia)
- Stomach pains; these may be a symptom of an inflamed pancreas (pancreatitis);
- Ringing in the ears (tinnitus)
- Reversible loss of hearing (usually associated with high doses or in patients with kidney problems)
- Various liver or gall-bladder problems, which can cause yellowing of the skin and/or eyes (jaundice) or pale stools with dark urine
- Chest pains
- Abnormal heart rhythms (including palpitations, a faster heartbeat, a life-threatening irregular heartbeat called torsades de pointes or abnormal ecg heart tracing) or heart stopping (cardiac arrest)
- Fever
- Anorexia
- Confusion
- Fits (seizures)
- Vertigo (problems with balance that can result in feelings of dizziness or sickness – particularly on standing)
- Hallucinations (seeing or hearing things that aren't there)
- Feeling generally unwell (malaise)
- Inflammation of the kidneys (a condition known as interstitial nephritis)
- Low blood pressure
- Serious skin rashes that may involve blistering and can cover large areas of the torso,



PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

**EROMYCIN 250 TABLET**  
**EROMYCIN 500 TABLET**  
Erythromycin 250mg Tablet  
Erythromycin 500mg Tablet

face and limbs (conditions known as Stevens Johnson syndrome, toxic epidermal necrolysis and erythema multiforme).

- Vomiting and irritability in young children between the age of 1 month and 12 months
- Visual impairment/blurred vision (Mitochondrial optic neuropathy).

**5. HOW TO STORE EROMYCIN TABLET**

- Keep all medicines out of the sight and reach of children
- Do not take this medicine after the expiry date which is stated on the label after “EXP”  
The expiry date refers to the last day of that month.
- Store below 30°C.
- Store in the original container.

**6. FURTHER INFORMATION.**



Each Eromycin 250 tablet contains Erythromycin 250 mg  
Each Eromycin 500 tablet contains Erythromycin 500mg

**What EROMYCIN TABLET looks like and contents of the pack**

1 X10 & 10 X10 tablets packed in blisters and placed in an inner carton with insert.

**SUPPLIER AND MANUFACTURER**

Fidson Healthcare Plc,  
Km.38, Lagos-Abeokuta Expressway,  
Sango-Ota, Ogun State, Nigeria  
234-(0)8077008888  
Customercare@fidson.com

Approval of PIL text	Prepared by	Approved by:	Approved by:
Signature		f 	
Date	09.10.2023	09.10.2023	
Designation	AMRA	HoM	HQRA